

# **Saving bones without risking brain - bisphosphonates and risk of stroke: matched case-control study**

## **Mini Abstract**

We investigated the association between bisphosphonate treatment and the risk of stroke using a large routine clinical dataset. We found no association between bisphosphonate treatment and risk of stroke, after adjusting for large number of clinical and demographic confounders.

## **Abstract**

### **Purpose**

There is conflicting evidence on the link between bisphosphonates and stroke with studies variously showing increased, decreased or unchanged risk. We investigated the association between bisphosphonate treatment and the risk of stroke using a large routine clinical dataset.

### **Methods**

We used a matched nested case-control study design analysing routinely collected electronic data from patients registered at primary care practices in England participating in the Royal College of General Practitioners Research and Surveillance Centre. Cases were patients aged 18 years or over, either living or dead, recorded as having had a stroke in the period 1<sup>st</sup> January 2005 to March 31<sup>st</sup> 2016. Each case was matched to one control according to age, sex, general practice attended and calendar time. Data were analysed using Stata, version 14.2. Conditional logistic regression was used to determine odds ratios for stroke according to bisphosphonate treatment and duration in cases compared with controls. We adjusted for disease risk groups, cardiovascular risk factors, treatments, smoking status, alcohol consumption, ethnicity, bisphosphonate types, fracture and socioeconomic status using IMD (Index of Multiple Deprivation).

### **Results**

We included 31,414 cases of stroke with an equal number of matched controls. Overall, 83.2% of cases and controls were aged 65 years or older and there were similar proportions of females (51.5%) and males (48.5%). Bisphosphonate treatment was not associated with stroke after adjusting for the wide range of confounders considered (OR 0.86, 95% CI 0.62 - 1.19).

1    **Conclusions**

2    We found no association between bisphosphonate treatment and risk of stroke, after adjusting for  
3    other confounders.

4

5

6    **Keywords**

7    Bisphosphonates, Stroke, Fracture, ONJ, Nested Matched Case Control study, Electronic Clinical-  
8    Patient Dataset.

9    **Manuscript word count: 4263**

## 1   **Introduction**

2   Screening, prevention and treatment of osteoporosis have increased in the UK, mainland Europe and  
3   the United States due to various factors including the need to address the increasing burden and costs  
4   of fractures, an ageing population at greater risk of fracture, increased awareness of risk factors for  
5   fracture and greater availability of screening tests (bone scans) and screening algorithms.[1]  
6

7   According to national guidance in the UK and US, bisphosphonate drugs (alendronate, pamidronate,  
8   risedronate, ibandronate, zoledronic acid) have largely replaced the use of Vitamin D with or without  
9   supplemental calcium for preventing fragility fractures NICE [2] in those deemed to be at-risk,[3, 4]  
10   mainly due to lack of evidence of effectiveness of the latter.[5, 6] Bisphosphonates prevent bone loss  
11   by slowing down the cells that break down and reabsorb old bone.  
12

13   There have also been conflicting systematic reviews about the association with cardiovascular disease  
14   (CVD) of treatments for osteoporosis. Neither calcium[7] nor vitamin D[8], which have been used for  
15   treatment of osteoporosis have been shown to be associated with cardiovascular benefits or adverse  
16   effects. Bisphosphonates, on the other hand, have shown conflicting evidence of CVD risk.  
17

18   Previous observational studies have suggested an increased risk, albeit small, of fatal stroke with  
19   bisphosphonates,[9] whereas other studies have not found such an association,[10] or found a  
20   reduction in risk of stroke with these agents.[11] Similar early reviews suggested that some  
21   bisphosphonates could be associated with atrial fibrillation, and although this is a heart rhythm  
22   disorder which in some cases can trigger stroke, these studies showed no association with stroke.[12,  
23   13] More recent reviews have shown no adverse effect on stroke, no reduction in cardiovascular  
24   outcomes overall,[14] but a modest increase in risk of atrial fibrillation.[15]  
25

26   We aimed to investigate the association between bisphosphonate treatment and stroke (fatal and non-  
27   fatal) using a large routine clinical dataset.  
28

## Methods

### *Study design*

We used a matched nested case-control study design to investigate the effect of bisphosphonates on risk of stroke. We identified cases (patients with stroke) and controls (patients without stroke) and compared these for prescriptions of bisphosphonates and other risk factors for stroke prior to the date of the stroke (or an equivalent date in control patients). Stroke was defined as both ischaemic and haemorrhagic stroke together with transient ischaemic attack or TIA, which is also known as a mini-stroke. TIA is the same as a stroke, except that the symptoms last for a short amount of time and no longer than 24 hours. This is because the blockage that stops the blood getting to the brain is temporary.

Key medications including bisphosphonates were defined and grouped using British National Formulary (BNF) chapter (Table 1).

1 Table 1 Drugs searched through BNF and defined through EMIS and Read Codes.

Bisphosphonates	BNF 6.6.2	Alendronic Acid, Etidronate, Ibandronic Acid, Pamidronate Disodium, Risedronate Sodium, Tilondronate, Sodium Clodronate, Zoledronic Acid.
Vasodilator antihypertensive drugs	BNF 2.5.1	Ambrisentan, Bosentan, Hydralazine Hydrochloride, Iloprost Macitentan, Minoxidil, Riociguat, Sildenafil, Sodium Nitroprusside, Tadalafil.
Centrally acting antihypertensive drugs	BNF 2.5.2	Clonidine Hydrochloride, Methyldopa, Moxonidine.
Adrenergic neurone blocking drugs	BNF 2.5.3	Guanethidine Monosulfate.
Alpha-adrenoceptor blocking drugs	BNF 2.5.4	Doxazosin , Indoramin, Prazosin, Terazosin, Phenoxybenzamine Hydrochloride, Phentolamine Mesilate
Angiotensin-converting enzyme inhibitors	BNF 2.5.5.1	Captopril, Enalapril, Maleate Fosinopril Sodium, Imidapril Hydrochloride, Lisinopril, Moexipril Hydrochloride, Perindopril, Erbumine, Perindopril, Arginine, Quinapril, Ramipril, Ramipril with Felodipine, Trandolapril.
Angiotensin-II receptor antagonists	BNF 2.5.5.2	Azilsartan, Candesartan, Eprosartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan.
Renin inhibitors	BNF 2.5.5.3	Aliskiren.
Calcium-channel blockers	BNF 2.6.2	Amlodipine, Diltiazem Hydrochloride, Felodipine, Isradipine, Lacidipine, Lercanidipine Hydrochloride, Nicardipine Hydrochloride, Nifedipine, Nimodipine, Verapamil Hydrochloride.
Statin	(Read & EMIS Codes)	Atorvastatin, Fluvastatin, Pravastatin Sodium, Rosuvastatin, Simvastatin, Simvastatin with Ezetimibe, Simvastatin with Fenofibrate.
Oral anticoagulant	(Read & EMIS Codes)	Warfarin Sodium, Acenocoumarol, Phenindione, Dabigatran, Etexilate, Rivaroxaban, Apixaban.

2  
3 We searched routinely collected electronic records from patients in England registered at primary care  
4 practices who are members of the Royal College of General Practitioners (RCGP) Research and  
5 Surveillance Centre (RSC).[16] This is a large computerised anonymised database representative of  
6 and comprising 2.9% of the population of England [17] including demographic information, data on  
7 health behaviours, referrals and treatment outcomes, with good clinical information including stroke  
8 and stroke deaths.[18] The study observational period was 1 January 2005 to 31 March 2016.

## *Selection of cases and controls*

The study cohort included patients drawn from all RCGP RSC practices over 10 years. Cases were patients aged 18 or over, either living or dead, recorded with standard computer codes for stroke. Each case of stroke was matched to one control according to age, sex, general practice attended and calendar time. Controls were patients registered at the same practice during the study period identified at the same index date as the corresponding case to account for possible seasonal effects and effects due to duration of observation for events. Controls were selected at random (and before their exposure status was known to reduce selection bias) from the pool of eligible matched controls for each case using incidence density sampling according to person-time at risk.[19] Controls had to be alive and not transferred out of the practice or dead prior to the index date of their matched case. All cases and controls that had less than 5 years of clinical records before the index date on the dataset were excluded to ensure completeness of recording of exposures and confounding variables. Those with a previous diagnosis of stroke were also excluded. For identified cases, the index date was the date of the first Stroke/TIA suffered by the patient. For the control, the index date was the date the patient they are matched to suffered their first stroke within the study period. The case was only stopped if it had deceased during the observational period.

## *Outcomes, exposures, confounder and effect modifiers*

Outcome measures were unadjusted and adjusted odds ratios and 95% confidence intervals for stroke associated with bisphosphonate treatment prior to the index date. Quintiles of the Index of Multiple Deprivation (IMD) were used as it ranks every small area in England from 1 to 32,844 (most deprived area to least deprived area) and Charlson Comorbidity Index (CCI) was used as a proxy for level of risk and frailty. [1] We adjusted for known confounding variables, in particular disease risk groups associated with stroke, cardiovascular risk factors, treatments, and other factors (ethnicity, Charlson comorbidity index, Index of Multiple Deprivation (IMD score) recorded with a computer (Read or EMIS) code. Fractures and bisphosphonate types were also included as confounding variables.

## *Data analysis*

Data were analysed using Stata, version 14.2 initially with some work done during the revision period using RStudio version 1.1.463. Descriptive statistics were given in terms of frequencies for categorical variables and means with standard deviations for continuous variables. We used conditional logistic regressions for matched case control studies, calculating unadjusted and adjusted odds ratios with 95% confidence intervals for stroke according to bisphosphonate treatment and duration in cases compared with controls. Adjusted analysis accounted for disease risk groups, cardiovascular risk factors, treatments, smoking status, alcohol consumption, fracture types, bisphosphonate types, ethnicity and socioeconomic status using the Index of Multiple Deprivation. All the confounding variables are listed in Tables (2-4). Fracture types, hip, osteonecrosis of the jaw (ONJ), vertebral and other fractures were grouped together. Bisphosphonate types were derived from Table 1.

### *Ethical approval*

We obtained ethical approval from Lincolnshire Community Health Services NHS Trust and the University of Lincoln, School of Health and Social Sciences Ethics Committee, the research protocol was approved by RCGP RSC.

## **Results**

### *Unadjusted analysis*

We included 31,414 cases of stroke with an equal number of matched controls (Table 2). Overall 83.2% in both cases and controls were aged 65 years or older and there were similar proportions of females and males (51.5% females and 48.5% males).

All the disease risk groups included in the analysis had a negative impact on the risk of stroke, with atrial fibrillation, peripheral arterial disease and hemiplegia being the groups with the highest impact on stroke (unadjusted OR of 2.13, 2.04 and 5.45 respectively; see Table 3). For cardiovascular risk factors (Table 2), being an ex-smoker, never smoker or non-drinker was associated with a reduced risk of stroke (unadjusted OR of 0.86, 0.82 and 0.82 respectively). The remaining cardiovascular risk

factors also had a negative impact on the risk of stroke, except high-density lipoprotein (HDL) cholesterol.

In relation to treatments (Table 4), we found that all were associated with a higher risk of stroke, with aspirin, statins and oral anticoagulants having the strongest association (unadjusted odds ratios of 1.79, 1.45 and 1.47, respectively). The unadjusted analysis also showed that a greater number of comorbidities were associated with a higher risk of stroke. Finally, the Index of Multiple Deprivation (IMD score) showed that people living in the most deprived areas had a higher risk of stroke than those living in the least deprived ones (without considering any other confounding).

The number of patients who had received bisphosphonates was greater for the cases than for controls: 9.3% of cases were prescribed bisphosphonates, while 7.6% of patients in the control group had received a prescription. The unadjusted odds ratio (OR) for the bisphosphonate treatment was 1.27, indicating that patients taking bisphosphonates were 27% more likely to suffer stroke than patients with no treatment, without considering any confounding variables.

Alendronate, Ibandronate and Risedronate showed reduced risk (unadjusted OR 0.78, 0.66, and 0.84 respectively) of association with stroke, and fracture types were not statistically significant for the unadjusted analysis. Most patients with lower CCI index took Alendronate followed by Risedronate and Ibandronate, Figure (1). Alendronate remained the preferred bisphosphonate of choice even as CCI index increased, showing highest prescription number for the first CCI index.






Figure 1. Showing number of bisphosphonate prescriptions uptake against Charlson Comorbidity Index.

#### Adjusted analysis

Table 5 shows the results of the adjusted analysis, where we included all the variables in the same regression to adjust for other confounders. The adjusted OR for the bisphosphonate treatment was 0.86 (95% CI 0.62 - 1.19) indicating the absence of a relationship between the drug and stroke, once all other confounding variables had been taken into account.

Among the disease risk groups, hemiplegia, atrial fibrillation and peripheral arterial disease were the variables with the highest effect on the risk of having stroke (OR 4.59, 1.98 and 1.48, respectively). Some of the disease risk groups included in the analysis, such as hyperlipidaemia, diabetes, chronic renal disease, Chronic Obstructive Pulmonary Disease (COPD), mild liver disease, peptic ulcer and rheumatological disease, showed a non-significant effect on the risk of stroke, when they were considered in the adjusted analysis.

Alcoholism (OR 1.43, 95% CI 1.30-1.57) and most deprived IMD (OR 1.27, 95% CI 1.18-1.37) were the risk factors showing the highest association with risk of stroke. Being a safe drinker was

associated with a lower risk of stroke (OR 0.84, 95% CI 0.80-0.88) compared to someone who had never drank alcohol, as were those on warfarin (OR 0.40, 95% CI 0.37-0.59).

The treatment associated with the highest risk of stroke was aspirin (OR 1.44, 95% CI 1.38-1.50), whereas antihypertensive treatment was associated with a reduced risk of stroke (OR 0.92). Finally, looking at IMD score, we found that patients living in the most deprived areas were more likely to suffer stroke compared to those living in the least deprived areas, as previously shown in the unadjusted analysis. The CCI showed a lower gradient than for the unadjusted analysis but still showed slight increase in odds ratio as number of comorbidities increased.

**Table 2 Characteristics of cases of stroke and matched controls**

Variables	Cases N=31,414, N (%)	Controls N=31,414, N (%)	Unadjusted OR (95% CI)	P-Value
Matching variables				
Age				
18-24 years	37 (0.1)	37 (0.1)	NA	
25-34 years	178 (0.6)	178 (0.6)		
35-44 years	404 (1.3)	404 (1.3)		
45-54 years	1561 (5)	1561 (5)		
55-64 years	3109 (9.9)	3109 (9.9)		
≥65 years	26125 (83.2)	26125 (83.2)		
Sex				
Female	16331 (52)	16331 (52)	NA	
Male	15083 (48)	15083 (48)		
Ethnicity				
White	21462 (68.3)	19004 (60.5)	ref	0.22
Asian	633 (2)	577 (1.8)	0.92 (0.81 - 1.05)	
Black	493 (1.6)	384 (1.2)	1.09 (0.94 - 1.27)	
Mixed	104 (0.3)	106 (0.3)	1.16 (0.87 - 1.56)	
Other	86 (0.3)	107 (0.3)	0.72 (0.54 - 0.97)	
Missing	8636 (27.5)	11236 (35.8)		
Fracture*				
Hip	355 (1.13%)	246 (0.78%)	ref	0.58
ONJ	10 (0.03%)	9 (0.03%)	1.30 (0.50 - 3.31)	
Other Fracture	2595 (8.26%)	1871 (5.96%)	1.04 (0.88 - 1.24)	
None	28322 (90.2%)	29193 (92.9%)	1.49 (1.26 - 1.75)	

<b>IMD quintile</b>				
1-Most deprived	4575 (14.6)	4114 (13.1)	ref	
2	4704 (15)	4449 (14.2)	1.35 (1.25 - 1.45)	p<0.001
3	6060 (19.3)	6059 (19.3)	1.19 (1.12 - 1.27)	p<0.001
4	7418 (23.6)	7381 (23.5)	1.16 (1.10 - 1.22)	p<0.001
5-Least deprived	8596 (27.4)	9116 (29.0)	1.07 (1.02 - 1.13)	p<0.001
Missing	61 (0.2)	295 (0.9)		

<b>Bisphosphonate Types<sup>‡*</sup></b>				
Alendronate	1975(6.29)	2488(7.92)	0.78 (0.73 - 0.83)	<0.001
none	29439(93.7)	28926(92.1)	ref	
Clodronate	23(0.07)	18(0.06)	1.28 (0.69 - 2.40)	0.441
none	31391(99.9)	31396(99.9)	ref	
Etidronate	101(0.32)	88(0.28)	1.15 (0.86 - 1.53)	0.345
none	31313(99.7)	31326(99.7)	ref	
Ibandronate	88(0.28)	133(0.42)	0.66 (0.50 - 0.86)	0.002
none	31326(99.7)	31281(99.6)	ref	
Risedronate	416(1.32)	494(1.57)	0.84 (0.74 - 0.96)	0.009

<sup>‡</sup> Some patients may have taken more than one type of Bisphosphonate over their prescription period. Only 1 person was prescribed Pamidronate so it is excluded in the table above.

<sup>\*</sup>Fracture and Bisphosphonate types added using RStudio.[20]

1 **Table 3 Disease risk groups and cardiovascular risk factors**

Variables	Cases N=31,414, N (%)	Controls N=31,414, N (%)	Unadjusted OR (95% CI)	P-Value
<b>Disease risk groups</b>				
CKD	6037 (19.2)	4670 (14.9)	1.42 (1.36 - 1.48)	p<0.001
COPD	2018 (6.4)	1615 (5.1)	1.28 (1.19 - 1.37)	p<0.001
Diabetes	4922 (15.7)	3644 (11.6)	1.43 (1.36 - 1.50)	p<0.001
Diabetes with complications	1635 (5.2)	973 (3.1)	1.74 (1.60 - 1.89)	p<0.001
Hyperlipidaemia	2934 (9.3)	2342 (7.5)	1.31 (1.24 - 1.39)	p<0.001
Acute myocardial infarction	1007 (3.2)	669 (2.1)	1.53 (1.39 - 1.69)	p<0.001
Angina	1555 (5.0)	1172 (3.7)	1.36 (1.26 - 1.48)	p<0.001
Atrial fibrillation	3290 (10.5)	1687 (5.4)	2.13 (2.00 - 2.27)	p<0.001
Congestive cardiac failure	1173 (3.7)	884 (2.8)	1.35 (1.23 - 1.48)	p<0.001
Hypertension	11134 (35.4)	8589 (27.3)	1.53 (1.47 - 1.58)	p<0.001
Peripheral arterial disease	771 (2.5)	389 (1.2)	2.04 (1.80 - 2.31)	p<0.001
Hemiplegia	258 (0.8)	36 (0.1)	5.45 (3.99 - 7.43)	p<0.001
Mild liver disease	257 (0.8)	176 (0.6)	1.47 (1.22 - 1.79)	p<0.001
Moderate liver disease	71 (0.2)	51 (0.2)	1.25 (0.88 - 1.77)	0.22
Peptic ulcer	474 (1.5)	294 (0.9)	1.59 (1.38 - 1.84)	p<0.001
Rheumatological disease	945 (3.0)	707 (2.3)	1.36 (1.23 - 1.5)	p<0.001
Cancer	3034 (9.7)	2870 (9.1)	1.07 (1.01 - 1.13)	0.02
Dementia	2041(6.5)	1440 (4.6)	1.51 (1.40 - 1.62)	p<0.001
<b>Cardiovascular risk factors</b>				
<b>Family history</b>				
Family history of stroke	1384 (4.4)	1054 (3.4)	1.34 (1.23 - 1.46)	p<0.001
Family history of ischemic heart disease	2109 (6.7)	1906 (6.1)	1.18 (1.10 - 1.26)	p<0.001
<b>Smoking</b>				
Active	699 (2.2)	431 (1.4)	ref	
Ex-Smoker	1898 (6)	1524 (4.9)	0.86 (0.75 - 0.99)	0.04
Never	1387 (4.4)	1224 (3.9)	0.82 (0.71 - 0.94)	0.01
Missing	27430 (87.3)	28235 (89.9)		
<b>Alcohol consumption category</b>				
Non-drinker	7333 (23.3)	6549 (20.9)	ref	
Safe	9580 (30.5)	10402 (33.1)	0.82 (0.79 - 0.87)	p<0.001
Hazardous	8935 (28.4)	8205 (26.1)	1.01 (0.95 - 1.06)	0.82
Alcoholism	1589 (5.1)	958 (3.1)	1.51 (1.37 - 1.66)	p<0.001
Missing	3977 (12.7)	5300 (16.9)		

2  
3  
4  
5  
6  
7  
8  
9

**Table 4 Clinical measurements, treatments (including bisphosphonates) and Charlson Index**

Variables	Cases N=31,414, N (%)	Controls N=31,414, N (%)	Unadjusted OR (95% CI)	P- Value
<b>Clinical measurements</b>				
Body Mass Index recorded	17973 (57.2)	16140 (51.4)		
Body Mass Index kg/m <sup>2</sup> (mean [SD])	27.3[5.5]	27.0[5.4]	1.01 (1.0 - 1.02)	p<0.001
Systolic blood pressure recorded	24171 (76.9)	22258 (70.9)		
Systolic blood pressure mmHg (mean [SD])	137.8 [19.1]	136.1 [18.1]	1.01 (1.01 - 1.01)	p<0.001
Diastolic blood pressure recorded	24171 (76.9)	22258 (70.9)		
Diastolic blood pressure mmHg (mean [SD])	77.6 [11.1]	76.8 [10.6]	1.02 (1.01 - 1.02)	0.04
LDL recorded	14061 (44.8)	11296 (36)		
LDL mmols/l (mean [SD])	2.9	2.8	1.04 (1.00 - 1.07)	p<0.001
HDL recorded	16682 (53.1)	13555 (43.1)		
HDL mmols/l (mean [SD])	1.5 [0.5]	1.5	0.90 (0.84 - 0.96)	p<0.001
Total cholesterol recorded	18836 (60.0)	15284 (48.7)		
Total cholesterol mmols/l (mean [SD])	5.0 [1.2]	5.0 [1.2]	1.04 (1.01 - 1.06)	p<0.001
Weekly alcohol units recorded	3370 (10.7)	3881 (12.4)		
Weekly alcohol units (mean [SD])	6.7 [13.6]	6.6 [12.8]	1.00 (1.00 - 1.01)	0.74
<b>Treatments</b>				
Aspirin uptake	11112 (35.4)	7660 (24.4)	1.79 (1.72 - 1.85)	p<0.001
Antihypertensive treatment	11963 (38.1)	9544 (30.4)	1.43 (1.38 - 1.48)	p<0.001
Statin uptake	10998 (35)	8768 (27.9)	1.45 (1.40 - 1.50)	p<0.001
Oral anticoagulant	2109 (6.7)	1655 (5.3)	1.3 (1.22 - 1.39)	p<0.001
Calcium uptake	10639 (33.9)	8447 (26.9)	1.47 (1.41 - 1.52)	p<0.001
Vitamin D uptake	4782 (15.2)	3830 (12.2)	1.35 (1.28 - 1.42)	p<0.001
Warfarin	1987 (6.3)	1615 (5.1)	1.25 (1.17 - 1.34)	p<0.001
<b>Bisphosphonates</b>				
Bisphosphonates uptake (last 3 years)	2909 (9.3)	2384 (7.6)	1.27 (1.19-1.34)	p<0.001
Number of bisphosphonate prescriptions (mean [SD])	24.0 [27.9]	22.2 [27.9]		
<b>Charlson Index</b>				
0	12116 (38.6)	15850 (50.5)	ref	
1	7170 (22.8)	6072 (19.3)	1.65 (1.58 - 1.73)	p<0.001
2	4612 (14.7)	3668 (11.7)	1.81 (1.71 - 1.91)	p<0.001
3	3454 (11)	2700 (8.6)	1.89 (1.78 - 2.01)	p<0.001
4	1917 (6.1)	1413 (4.5)	2.09 (1.94 - 2.26)	p<0.001
≥5	2145 (6.8)	1711 (5.5)	2.05 (1.92 - 2.19)	p<0.001

1 **Table 5 Conditional logistic regression for matched cases and controls model adjusted for**  
2 **confounders**

Stroke/TIA		Odds Ratio	[95% Conf.	Interval]	P-value
Bisphosphonates Types	All Bisphosphonates	0.86	0.62	1.19	0.35
	Alendronate	1.13	0.81	1.57	0.46
	Clodronate	0.98	0.46	2.08	0.96
	Etidronate	0.89	0.57	1.40	0.61
	Ibandronate	1.31	0.83	2.06	0.24
	Risedronate	1.12	0.79	1.58	0.54
	ZoledronicAcid	0.79	0.21	2.95	0.73
Drugs	Calcium	0.94	0.9	0.99	0.03
	VitaminD	0.94	0.87	1.01	0.07
	Aspirin	1.44	1.38	1.50	p<0.001
	Antihypertensive	0.92	0.88	0.97	p<0.001
	Statins	0.99	0.95	1.04	0.83
	Warfarin	0.40	0.27	0.59	p<0.001
	Oral AntiCoagulant Therapy	1.87	1.27	2.75	p<0.001
Diseases	Chronic kidney disease	0.95	0.88	1.01	0.1
	Chronic Obstructive Pulmonary Disease	0.97	0.91	1.06	0.71
	Diabetes	0.97	0.91	1.03	0.3
	Acute myocardial infarction	1.13	1.01	1.25	0.03
	Angina	0.97	0.89	1.05	0.44
	Atrial fibrillation	1.98	1.83	2.13	p<0.001
	Congestive cardiac failure	0.91	0.82	1.01	0.07
	Hypertension	1.11	1.05	1.17	p<0.001
	Peripheral arterial disease	1.48	1.30	1.69	p<0.001
	Hemiplegia	4.59	3.33	6.34	p<0.001
	Immunosuppression	0.98	0.83	1.16	0.79
	Mild liver disease	0.95	0.77	1.17	0.65
	Moderate liver disease	0.73	0.50	1.07	0.11
	Peptic ulcer	1.25	1.07	1.46	p<0.001
	Rheumatological disease	1.11	0.99	1.24	0.07
	Hyperlipidaemia	1.09	1.02	1.16	0.01
Life Style	Active smoker	1.15	0.99	1.33	0.07
	Ex-smoker	1.00	0.90	1.11	0.98
	Safe alcohol consumption	0.84	0.80	0.88	p<0.001
	Hazardous alcohol consumption	0.96	0.92	1.02	0.18
	Alcoholism	1.43	1.30	1.57	p<0.001
Fracture	Hip	1.39	1.17	1.66	p<0.001
	ONJ	1.09	0.43	2.75	p0.85
	Other Fractures	1.35	1.26	1.45	p<0.001
	Vertebral	1.33	1.01	1.76	0.05

**Deprivation Quintiles**

DeprivationQuintile1	<b>1.27</b>	1.18	1.37	p<0.001
DeprivationQuintile2	<b>1.15</b>	1.07	1.22	p<0.001
DeprivationQuintile3	<b>1.13</b>	1.07	1.2	p<0.001
DeprivationQuintile4	<b>1.06</b>	1.01	1.12	0.01

**Co-morbidity**

CharlsonI1	<b>1.37</b>	1.30	1.45	p<0.001
CharlsonI2	<b>1.45</b>	1.35	1.55	p<0.001
CharlsonI3	<b>1.41</b>	1.29	1.54	p<0.001
CharlsonI4	<b>1.45</b>	1.29	1.62	p<0.001
CharlsonI5	<b>1.38</b>	1.21	1.56	p<0.001

---

Analysis performed using R package Survival. [21]

## Discussion

### *Main findings*

We found no association between bisphosphonate treatment and risk of stroke, after adjusting for age, sex, ethnicity, index of multiple deprivation, Charlson index, Fracture status, clinical and treatment variables.

### *Strengths and limitations*

The limitations of the case-control approach include confounding and residual unmeasured confounding. The large validated database we used enabled us to adjust for important confounders including clinical risk groups, cardiovascular risk factors and differences in treatment between cases and controls. We also adjusted for additional comorbidities using the Charlson index. Although great care was taken to ensure all stroke/TIAs were correctly coded, around 20% of all stroke patients' are those who have temporary symptoms of TIA episodes.[22] These are difficult to diagnose as they depend on patient history. Because the duration of the episode is short, patients' symptoms are likely to have resolved by the time of assessment, and the absence of an established biomarker makes the diagnosis difficult. Information about physical activity and diet were not available and these factors may be a source of residual confounding.

### *Comparison with previous studies*

The previous contradictory evidence linking bisphosphonates and stroke provided the rationale for this study. Although bisphosphonates have been associated with atrial fibrillation, presumed to be an idiosyncratic adverse effect which can sometimes lead to an embolic stroke due to thrombus generated in the abnormally contracting atrium blocking a narrowed carotid artery, there was no association with stroke in these studies.[12, 13]

Other studies have suggested that bisphosphonates may prevent cardiovascular disease including stroke and myocardial infarction mediated through a reduction in vascular calcification or reduced atherosclerosis through a number of mechanisms.[23] Vestergaard and colleagues, in a large cohort



1 study in Denmark found a reduction in overall risk of cardiovascular events but an increase in fatal  
2 strokes but the variability in effect for different drugs and the differences in risk observed were  
3 small.[9]

4  
5 One study showed no association between prior bisphosphonate therapy and 30-day mortality from  
6 stroke.[24] The most recently published meta-analysis, from Kim and colleagues, concluded that  
7 ‘bisphosphonates do not have beneficial or harmful effects on atherosclerotic cardiovascular events,  
8 but zoledronic acid may modestly increase the risk of atrial fibrillation’.[15]

#### 9 10 *Bisphosphonate Compliance*

11 There is a question about those patients who are possibly at a higher risk may have been less likely to  
12 take the bisphosphonates, figure (1). Generally, there is a problem with compliance as Park and  
13 colleagues have shown that compliance and persistence with oral bisphosphates in patients with  
14 rheumatoid arthritis were suboptimal in real practice, thereby limiting the efficacy of osteoporosis  
15 treatment. [25]This may be true for our study as well.

#### 16 17 *Implications for policy, practice and research*

18 This study suggests that bisphosphonate as a group of drugs are not associated with increased risk of  
19 stroke. This and evidence of effectiveness for prevention of osteoporosis supports their use first-line  
20 for prevention of osteoporosis in those deemed to be at high risk.[2] Clinicians will continue to use  
21 these drugs but greater consideration is being given to the time to stop [26] because of a lack of  
22 benefit beyond this time. There is limited evidence that bisphosphonates can cause painful, hard-to-  
23 treat osteonecrosis damage to the jaw bone, as well as very rare fractures of the mid-femur, further  
24 research is needed in these areas to shed more light. Our study finds no evidence for excess ONJ  
25 fracture due to bisphosphonates however there were noticeably more hip and other fractures present in  
26 the cases. Finally, we recommend further research to incorporate other confounders, to conduct  
27 studies which overcome unknown or unmeasured confounders for example using self-controlled case  
28 series designs.

## Conclusion

We found no association between bisphosphonate treatment and risk of stroke, after adjusting for other confounders. Bisphosphonates are considered first-line for prevention of osteoporosis and fragility fractures and this study supports their safety in people at risk of stroke.

## References

1. Quan, H., et al., *Updating and validating the Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using data from 6 countries*. Am J Epidemiol, 2011. **173**(6): p. 676-82.
2. NICE, *Bisphosphonates for treating osteoporosis*. 2017, London: National Institute for Health and Clinical Excellence.
3. NICE, *Osteoporosis: assessing the risk of fragility fracture*. 2012, London: National Institute for Health and Clinical Excellence.
4. Qaseem, A., et al., *Treatment of low bone density or osteoporosis to prevent fractures in men and women: a clinical practice guideline update From the American College of Physicians*. Ann Intern Med, 2017. **166**(11): p. 818-839.
5. Reid, I.R., M.J. Bolland, and A. Grey, *Effects of vitamin D supplements on bone mineral density: a systematic review and meta-analysis*. Lancet, 2014. **383**(9912): p. 146-55.
6. Kahwati, L.C., et al., *Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-dwelling adults: evidence report and systematic review for the US Preventive Services Task Force*. JAMA, 2018. **319**(15): p. 1600-1612.
7. Chung, M., et al., *Calcium intake and cardiovascular disease risk: an updated systematic review and meta-analysis*. Ann Intern Med, 2016. **165**(12): p. 856-866.
8. Ford, J.A., et al., *Cardiovascular disease and vitamin D supplementation: trial analysis, systematic review, and meta-analysis*. Am J Clin Nutr, 2014. **100**(3): p. 746-55.
9. Vestergaard, P., et al., *Stroke in relation to use of raloxifene and other drugs against osteoporosis*. Osteoporos Int, 2011. **22**(4): p. 1037-45.

- 1 10. Christensen, S., et al., *Oral bisphosphonates and risk of ischemic stroke: a case-control study*.  
2 Osteoporos Int, 2011. **22**(6): p. 1773-9.
- 3 11. Kang, J.H., J.J. Keller, and H.C. Lin, *A population-based 2-year follow-up study on the*  
4 *relationship between bisphosphonates and the risk of stroke*. Osteoporos Int, 2012. **23**(10):  
5 p. 2551-7.
- 6 12. Sharma, A., et al., *Risk of atrial fibrillation with use of oral and intravenous bisphosphonates*.  
7 Am J Cardiol, 2014. **113**(11): p. 1815-21.
- 8 13. Sharma, A., et al., *Risk of serious atrial fibrillation and stroke with use of bisphosphonates:*  
9 *evidence from a meta-analysis*. Chest, 2013. **144**(4): p. 1311-1322.
- 10 14. Kranenburg, G., et al., *Bisphosphonates for cardiovascular risk reduction: A systematic review*  
11 *and meta-analysis*. Atherosclerosis, 2016. **252**: p. 106-115.
- 12 15. Kim, D.H., et al., *Bisphosphonates and risk of cardiovascular events: a meta-analysis*. PLoS  
13 One, 2015. **10**(4): p. e0122646.
- 14 16. Correa, A., et al., *Royal College of General Practitioners Research and Surveillance Centre*  
15 *(RCGP RSC) sentinel network: a cohort profile*. BMJ Open, 2016. **6**(4): p. e011092.
- 16 17. de Lusignan, S., et al., *RCGP Research and Surveillance Centre Annual Report 2014-2015:*  
17 *disparities in presentations to primary care*. Br J Gen Pract, 2017. **67**(654): p. e29-e40.
- 18 18. Hinton, W., et al., *Incidence and prevalence of cardiovascular disease in English primary care:*  
19 *a cross-sectional and follow-up study of the Royal College of General Practitioners (RCGP)*  
20 *Research and Surveillance Centre (RSC)*. BMJ Open, 2018. **8**(8): p. e020282.
- 21 19. Knol, M.J., et al., *What do case-control studies estimate? Survey of methods and assumptions*  
22 *in published case-control research*. Am J Epidemiol, 2008. **168**(9): p. 1073-81.
- 23 20. Team, R., *RStudio: Integrated Development for R*. RStudio, B. Inc., MA URL, Editor. 2018.
- 24 21. T, T., *A Package for Survival Analysis in S\_*. version 2.38. 2015.
- 25 22. Fitzpatrick, T., et al., *How do neurologists diagnose transient ischemic attack: A systematic*  
26 *review*. International Journal of Stroke. **0**(0): p. 1747493018816430.

- 1 23. Caffarelli, C., et al., *Bisphosphonates, atherosclerosis and vascular calcification: update and*  
2 *systematic review of clinical studies*. Clin Interv Aging, 2017. **12**: p. 1819-1828.
- 3 24. Christensen, D.H., et al., *The impact of preadmission oral bisphosphonate use on 30-day*  
4 *mortality following stroke: a population-based cohort study of 100,043 patients*. Clin  
5 Epidemiol, 2015. **7**: p. 381-9.
- 6 25. Park, J.-H., et al., *Compliance and persistence with oral bisphosphonates for the treatment of*  
7 *osteoporosis in female patients with rheumatoid arthritis*. BMC musculoskeletal disorders,  
8 2017. **18**(1): p. 152-152.
- 9 26. Paskins, Z. and L. Warburton, *Bisphosphonates beyond five years*. BMJ, 2016. **352**: p. i264.

10

11